

Resonant tunneling through a single level quantum dot attached to ferromagnetic leads with non-collinear magnetizations

M. Wawrzyniak¹ and J. Barnaś^{1,2}

¹*Department of Physics, Adam Mickiewicz University
Umultowska 85, 61-614 Poznań, Poland*

²*Institute of Molecular Physics, Polish Academy of Sciences
M. Smoluchowskiego 17, 60-179 Poznań, Poland*

Resonant electronic transport through a non-interacting single-level quantum dot attached to ferromagnetic leads is analyzed theoretically. Magnetic moments of the leads are assumed to be non-collinear. Apart from this, an external magnetic field, non-collinear with the magnetizations, is applied to the system. However, magnetic moments of the leads and the external magnetic field are in a common plane. Basic transport characteristics, including current voltage curves, linear and non-linear conductance, and tunnel magnetoresistance associated with magnetization rotation are calculated using the non-equilibrium Green function technique, and the Green functions has been calculated via the equation of motion method. Dependence of the transport characteristics on the system parameters, like energy level position, spin polarization of the leads, asymmetry between right and left leads, has been calculated numerically and the results are discussed from the fundamental and application point of view.

Name of the presenting author (poster): Małgorzata Wawrzyniak
e-mail address: lauraphy@spin.amu.edu.pl
url's: <http://www.fizyka.amu.edu.pl>